

Series: The Biology of Antigen Presentation

Feature Review

Present Yourself! By MHC Class I and MHC Class II Molecules

Kenneth L. Rock,¹ Eric Reits,² and Jacques Neefjes^{3,4,*}

Since the discovery of MHC molecules, it has taken 40 years to arrive at a coherent picture of how MHC class I and MHC class II molecules really work. This is a story of the proteases and MHC-like chaperones that support the MHC class I and II molecules in presenting peptides to the immune system. We now understand that the MHC system shapes both the repertoire of presented peptides and the subsequent T cell response, with important implications ranging from transplant rejection to tumor immunotherapies. Here we present an illustrated review of the ins and outs of MHC class I and MHC class II antigen presentation.

Why Do We Need to Present Antigens?

T cells help eliminate pathogens present in infected cells and also help B cells make better and different kinds of antibodies to protect against extracellular microbes and toxic molecules. To accomplish these important functions, T cells have to interact intimately with other cells and then find and instruct or eliminate the ones that are harboring or have been exposed to these pathogenic threats. However, T cells are unable to peek beneath the surface of cells to identify ones that have ingested bacteria or are synthesizing viral or mutant proteins. Instead, antigen presentation systems have evolved that display on the cell surface information about the various antigens that cells are synthesizing or have ingested. These antigen presentation pathways monitor the major subcellular compartments where pathogens could be lurking and report their findings to the appropriate kinds of T cells. Endogenously synthesized antigens in the cytosol of all cells are presented to CD8⁺ T cells as peptides bound to MHC I molecules, thereby allowing the CD8⁺ lymphocytes to identify and eliminate virally infected cells or cancers. Antigens ingested into endocytic compartments of macrophages, dendritic cells, or B cells are presented to CD4⁺ T cells as peptides bound to MHC II molecules. T cells have antigen receptors that recognize antigenic peptide, but only in the context of MHC I or MHC II molecules that are displaying the antigen on the cell surface. Consequently, T cells are directed to work with cells while not being 'distracted' by free antigen, against which they would be unable to do anything. Moreover, the pattern of expression of MHC I and II molecules directs T cells to interact with exactly the right kind of cells. Here we review the current understanding of the mechanisms of antigen presentation as well as their implications in health and disease. The studies discussed here have paved the way for the increasingly refined analyses of the biology of antigen presentation – in various physiological and clinical contexts, cells, and organs – that are the focus of this special issue.

The MHC Scaffold, or How Did MHC I and MHC II Molecules Evolve?

MHC I and II molecules present protein fragments to CD8⁺ and CD4⁺ T cells, respectively. These molecules are essential for cell-mediated immunity and therefore appeared at the inception of the adaptive immune system 500 million years ago [1]. For their construction they used two

Trends

MHC molecules are critical in transplantation, autoimmunity, infections, and tumor immunotherapy.

The biology of antigen presentation by MHC I and MHC II molecules provides targets for manipulation of these diseases.

This biology also explains the fragments presented to the immune system and the cellular evolutionary race to escape immune control of infected and transformed cells.

Many of the players determining antigen degradation and subsequent peptide loading on MHC molecules are defined and are helping to improve the accuracy of predicting presented fragments.


The combined understanding of antigen presentation by MHC molecules allows exploitation to improve the responses of the cellular arm of the immune system to vaccinations and immunotherapies.

¹Department of Pathology, UMass Medical School, Worcester, MA, USA

²Department of Cell Biology and Histology, Amsterdam Medical Center, University of Amsterdam, Amsterdam, The Netherlands

³Department of Cell Biology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

⁴Department of Chemical Immunology, Leiden University Medical Center, Leiden, The Netherlands



*Correspondence:
J.J.C.Neeffjes@lumc.nl (J. Neeffjes).

Trends in Immunology


Figure 1. MHC Locus and Allelic Products with Polymorphism. Most mammalian species express three different MHC I and three different MHC II molecules (shown here for the human MHC I HLA-A, HLA-B, and HLA-C locus products). Since these are polymorphic and genetically encoded, between three and six (depending on differences between the inherited genes) alleles will be expressed on cells. These are polymorphic in the peptide-binding groove region of MHC molecules to present different peptides of a defined antigen to a CD8⁺ T cell (CTL).

immunoglobulin domains topped by two parallel alpha helices resting on a platform of beta-pleated sheets. This capital structure generated a peptide-binding groove between the alpha helices [2,3] that is, 'evolutionarily speaking', likely to have been borrowed from earlier chaperone structures. There are arguments for this: (i) chaperones bind unfolded stretches of proteins and the prototypic unfolded structure is a peptide; (ii) various phylogenetically older chaperones have a somewhat similar structure [4]; and (iii) there are various MHC molecule 'look alikes' that act as specific chaperones in the process of MHC-restricted antigen presentation. These include tapasin [5], tapasin-binding protein related (TAPBPR) [6,7], DM [8,9], and DO [10], as we discuss below. However, MHC I and MHC II molecules are unique in the proteome because of their extreme polymorphism (>10 000 different alleles of MHC I molecules have been identified thus far). This has interesting consequences. Polymorphic residues on the top alpha helices interact with the T cell receptor (TCR) and are the basis for the specificity of TCRs for an antigen peptide plus a particular allelic form of an MHC molecule (a phenomenon called MHC restriction). Polymorphic residues in the MHC peptide-binding groove change the nature and location of so-called pockets. These variable pockets are filled by complementary variable amino acid side chains of peptides (so-called anchor residues), with the effect that different fragments from a defined antigen are presented by different polymorphic MHC molecules [11,12] (Figure 1). Next to the anchor residues, most other amino acids in a peptide fill a free space and can be (almost) any of the 20 amino acids [13,14]. By having pockets with specificity for only a few side chains and allowing the remaining six to ten amino acids to vary between all possibilities, each kind of MHC molecule can present a very large repertoire of peptides. Moreover, by having three to six different MHC I as well as three to 12 different MHC II molecules (the exact number depending on how many different MHC alleles were inherited from one's parents and how the MHC II subunits are paired), cells can present a large fraction of the universe of peptides, although not all sequences. In theory, then, MHC I molecules can present a peptidome of around $6 \times 10^{6-7}$

different peptides and MHC II can display up to 12×10^{10} . In actuality, such a large array of peptides cannot all be presented because there are only around 200 000 MHC I and 20 000 MHC II molecules on cells such as B and T cells [15]. Moreover, since some peptides are present in high numbers (from highly expressed proteins), the real number of different peptides presented by one cell is likely to be less than 10 000. Importantly, when a pathogen alters a critical anchor residue in one of its antigenic epitopes, it may prevent presentation of this antigen in one individual but not in another with different MHC molecules that will simply select different peptides from the same pathogen [16]. Therefore, MHC polymorphism is good for the survival of the population and not necessarily the individual. How is this extensive polymorphism maintained in the species? One possibility is based on evidence that females can distinguish by smell MHC allele differences in males and prefer as mates individuals with whom they do not share MHC alleles. The consequence of such a preference would be to promote maximal expression of polymorphic MHC alleles in offspring and the maintenance of diverse alleles in the population [17]. In the human species, perfumes may mask the natural scent of this basis of partner choice, with unknown effects. The obvious modern disadvantage of MHC polymorphism is transplant rejection, but even this may serve a useful function in nature by preventing the seeding of cancer cells between individuals. This is illustrated by the fact that MHC-deficient oral cancers are currently being transferred between Tasmanian devils through bites and decimating the population of these animals in the wild [18,19].

How to Present Your Inner Self? MHC Class I Molecules

MHC I molecules present peptides from the proteins that are synthesized by cells. In healthy cells, all of these proteins are autologous ones to which CD8⁺ T cells are tolerant. However, when cells are expressing mutant sequences (e.g., in cancers), microbial genes (e.g., from viral infections), or foreign polymorphic genes (e.g., from transplants), these 'non-self' antigenic peptides are included in the presented peptidome, allowing CD8⁺ T cells to detect and destroy these abnormal cells. How MHC I manages to present a blueprint of the intracellular proteome has been established over the past 25 years [20] (Figure 2). Briefly, normal and abnormal proteins are degraded by the proteasome into peptide fragments [21,22]. These fragments are further trimmed and to a large extent destroyed by cytosolic peptidases [23] but some survive by escaping into the endoplasmic reticulum (ER) through a peptide transporter called transporter associated with antigen processing (TAP) that is embedded in the ER membrane [20]. In the ER, TAP forms the center of a peptide-loading complex that includes a dedicated chaperone, tapasin, empty MHC I molecules awaiting peptides, and two common chaperones, calreticulin and the protein disulfide isomerase ERp57 [24]. TAP translocates peptides that then are considered for binding by MHC I molecules. These empty MHC I molecules are held in a peptide-receptive state by the chaperone tapasin and tapasin also promotes MHC I binding of peptides with a slow off rate, thereby helping to shape the repertoire of presented peptides [25]. In this reaction, MHC I molecules test the binding of many peptides and subsequently release most of these until a proper (low-off-rate) peptide is bound [26]. These are usually peptides of a very specific length of eight to ten amino acids with appropriate anchor residues. Peptides that are too long can be trimmed by an ER-resident aminopeptidase, ERAP1 (and ERAP 2 in some species) [27–29], before consideration by MHC I molecules that are either in the peptide-loading complex or associating with another tapasin look-alike chaperone in the ER called TAPBPR. Like tapasin, TAPBPR also shapes the peptide repertoire on MHC I molecules [30,31]. Interestingly, binding of peptides longer than eight or nine residues, but not shorter ones, triggers a conformational change in ERAP1 that activates its hydrolysis [32–34]. Through this mechanism, ERAP1 trims most peptides down to eight or nine residues, corresponding to the size needed for optimal binding to MHC I molecules. Ultimately, the peptides that are available to be presented are the ones that have been cleaved to the right size and have somehow escaped further hydrolysis to a size that is too small to stably bind to MHC I molecules (Figure 3). Peptides that are unable to bind an MHC I molecule are ultimately translocated back into the cytosol for



Trends in Immunology


Figure 2. A Simple Illustration of MHC I Antigen Presentation. Antigens are degraded by the proteasome to yield peptide fragments. These peptides are then translocated from the cytosol into the endoplasmic reticulum (ER) lumen where MHC I waiting for peptides is retained by a series of chaperones including the dedicated chaperone tapasin in the peptide-loading complex. A second dedicated chaperone [tapasin-binding protein related (TAPBPR)] can further optimize the peptides in MHC I. Only MHC I with optimal peptides is allowed to leave the ER to present the peptide fragments at the cell surface to CD8⁺ T cells (CTL).

degradation [35]. Whether there are mechanisms that help to protect some of these peptides from destruction or release from the ER during the time before they bind MHC I molecules is not entirely clear.

Peptides should be considered the third subunit of MHC I molecules as they are required to stabilize these complexes when the MHC I molecules are not bound to chaperones in the ER [36]. Peptides also allow the MHC I molecules to be released from the ER quality control system (the various chaperones) for transport to the cell surface for presentation to CD8⁺ T cells [37,38]. This system of low-off-rate peptide selection, exporting only peptide-loaded MHC I complexes, may help prevent healthy cells from easily replacing their bound endogenous peptides for exogenous antigenic peptides, an event that would lead to the presentation of peptides that do not reflect the status of a given cell and possible execution by CD8⁺ T cells.

Complexity in the MHC Class I Antigen Presentation Pathway

The general scheme outlined above is somewhat more complex when taking into consideration the diversity of the MHC I family. The different loci expressed (in humans, HLA-A, -B, and -C) and the many polymorphic allelic forms behave somewhat differently with respect to expression, peptide binding, and stability [39,40]. Peptides may also be generated from multiple sources. This is because the proteasome not only degrades proteins as part of normal protein turnover but also degrades abnormal ones that arise from errors in translation, folding, and/or pairing. The degradation of these disabled proteins prevents their aggregation and also potentially more directly couples protein translation to antigen presentation [41–43]. Such antigens are called defective ribosomal products (DRiPs) and may be produced in greater amounts under high-protein-synthesis-conditions, such as occur during viral infection. The degradation of DRiPs



Trends in Immunology

Figure 3. Survival of the Fittest for MHC I Presentation and the Many Proteasomes. There are three types of proteasome with unique tissue expression. These proteasomes have altered cleavage specificities yielding (in part) different degradation fragments. These fragments are released in the hostile environment of the cytosol where most of the peptides will be destroyed by peptidases. A few peptides survive this massacre through translocation in the ER by transporter associated with antigen processing (TAP). Here they can be further trimmed by ER-resident aminopeptidase (ERAP) or translocated back into the cytosol by the ER-associated degradation (ERAD) system. Only a few peptides survive chaperone-mediated survival selection for low-off-rate peptides for a defined MHC I allele and these are ultimately presented to a CD8⁺ T cell (CTL).

would quickly generate peptides after initial translation of the antigen and this may allow rapid detection of infected cells [44,45].

While the source of antigens can vary, so can the cleavage of these antigens by different proteasomes. Many presented peptides can be generated through the phylogenetically older 'conventional' proteasome. However, the development of the immune system coincided with the evolution of alternative forms of active-site subunits for this particle, leading to the assembly of an immunoproteasome. A set of these subunits ($\beta 1i$, $\beta 2i$, and $\beta 5i$) [46] is constitutively expressed in dendritic cells and lymphocytes and can be induced in all other cells by interferons; for example, during viral infections [47,48]. When these subunits are expressed, they preferentially incorporate into newly assembling particles to form immunoproteasomes that generate a distinct set of peptides during protein degradation [49]. This shift from constitutive to immunoproteasome in cells often enhances the generation of peptides presented by MHC I molecules, including many unique ones [50]. Generation of new peptides will be at the cost of other peptide fragments that are cleaved to make the new fragment [51]. Another set of alternative active-site subunits ($\beta 1i$, $\beta 2i$, and $\beta 5t$) is expressed uniquely only in cortical thymic epithelial cells (cTECs), where they are incorporated into thymoproteasome particles [52]. Among the peptides generated by thymoproteasomes, several are unique and these play a critical role in the auditioning of developing CD8⁺ T cells during positive selection and in allowing many of these cells to avoid subsequent negative selection [50]. Proteasomes, but also the cytosolic and ER-associated peptidases, are variable in content and number. From this stew of proteolytic activities an

Not How to Present, But What to Present

How to Hide One's Inner Self? Pathogen and Tumor Escape from Antigen Presentation

Figure 4. Various Viral Immune Evasion Strategies for MHC I Antigen Presentation. Pathogens have evolved different ways to obstruct processes selective in the antigen presentation pathway. Examples include viral proteins inhibiting peptide transport by transporter associated with antigen processing (TAP), retaining MHC I in the endoplasmic reticulum (ER), and recognizing MHC I in the ER to target these back into the cytosol for degradation by the proteasome [the ER-associated degradation (ERAD) system]. Other viral proteins recognize MHC I at the cell surface for internalization and destruction in lysosomes.


ways to tamper with the process of antigen presentation [62]. CMV encodes proteins that inhibit the peptide transporter TAP or that induce the degradation of MHC I molecules in the ER or plasma membrane [63,64]. Other viruses shut down genomic MHC class I expression [65], produce a peptide mimic that blocks TAP, or block the transport of MHC I molecules to the cell surface [66]. Any step in the MHC class I antigen presentation pathway not interfering with cell viability can be expected to be manipulated by viruses to prevent their presentation.

Since most of the steps in the MHC class I pathway are not essential for viability and because cancer cells are often genetically unstable, tumors can, under the selection pressure imposed by CD8⁺ T cells, generate variants that have lost key components of the MHC class I pathway and escape control by CD8⁺ T cells [67]. Reduction of MHC I molecule expression in human tumors is often detected by pathologists [68]. This is one of the reasons for assuming an important role of the immune system in the control of particular tumors, especially those with more neoantigens (from mutated proteins) such as melanoma and lung cancer [69,70]. Understanding how a given tumor can evade detection by CD8⁺ T cells could help in determining the immunotherapies that are most likely to succeed against that tumor.

Some Exceptions in Self-Presentation by MHC Class I Molecules

The MHC class I antigen presentation system is constructed in such a way that most cells exclusively present their own antigens. Because of this, it was thought that CD8⁺ T cells selectively eliminate infected cells without destroying neighboring 'innocent bystanders'. However, this concept is challenged by the fact that the cytosols of many healthy cells (unlike cancer cells) are connected by so-called gap junctions. These gap junctions allow peptide fragments to pass into directly connected neighboring cells for entry in their antigen presentation pathway and presentation through the neighbor's MHC I molecules to CD8⁺ T cells [71,72]. It is likely that cytosolic peptidases will limit the spread of such peptides beyond the most proximal neighboring cells. However, since the proximal cells are at high risk of viral invasion, gap junctions may allow their elimination even before real entry of the pathogen.

A more intensely studied system where MHC I presents antigens that differ from those made by the cell itself is cross-presentation [73,74] (Figure 5). This pathway operates in dendritic cells and other phagocytes and is important because it plays a central role in immune surveillance [74–76]. It allows dendritic cells to acquire antigens from other infected cells and cancers in the periphery and then report their presence to naïve CD8⁺ T cells in lymphoid organs in ways that initiate an immune response. Phagocytes acquire these antigens when they ingest them by phagocytosis (e.g., by eating cell debris [77–79] or possibly even taking a 'bite' of living cells [80]) or via receptor-mediated endocytosis (e.g., of glycan-modified proteins through lectin receptors or antibody-bound antigens through Fc receptors [81,82]). There is more than one mechanism by which ingested antigens can be cross-presented, but one is likely to involve cytosolic transfer of antigen from the phagosome into the cytosol for degradation by the proteasome [83,84]. The fragments may then be loaded on MHC I molecules in the ER or translocated back into the phagosome for local MHC class I peptide loading [78]. In another mechanism, some antigens are degraded by lysosomal proteases and loaded onto recycling MHC class I molecules in a pathway similar to MHC class II molecules [85–87]. Toll-like receptor (TLR) signaling can induce the accumulation of MHC I in recycling endosomes to promote cross-presentation [88]. A potential limitation of this latter mechanism is that it could result in priming CD8⁺ T cells to peptides differing from the ones that the T cells will encounter in infected cells (as the priming peptides are generated by proteases different from the proteasome). However, this may also be a problem in the phagosome-to-cytosol mechanism of cross-presentation because, in the absence of inflammation, immunoproteasomes in dendritic cells may generate different fragments compared with the constitutive proteasome in most peripheral (and, notably, tumor) cells [89]. Thus, whether and how cross-presentation allows activation of CD8⁺ T cells recognizing




Trends in Immunology

Figure 5. Crossing Boundaries: Various Models of Cross-Presentation by MHC I. MHC I molecules can present exogenous antigens and antigens delivered in apoptotic bodies and other types of cell debris. For dendritic cells, this process can result in cross-priming of CD8⁺ T cells. MHC I can recycle through the endosomal pathway to acquire antigen fragments made by proteases such as insulin-regulated aminopeptidase (IRAP) [129] for cross-presentation. Endosomes may also acquire transporter associated with antigen processing (TAP) and other endoplasmic reticulum (ER) molecules that may help export antigens into the cytosol for proteasomal hydrolysis; the resulting peptides may be re-imported into the endosomes for MHC I loading and recycling and/or be delivered in the normal antigen presentation pathway as shown in Figure 2.

the full range of antigenic peptides presented by peripheral cells is unclear. Regardless of the exact mechanism, cross-presentation allows MHC I molecules to present peptides from antigens that usually are handled by MHC II molecules. The system employed by MHC II molecules for antigen presentation is understood in considerably more molecular detail and may share elements used for cross-presentation by MHC I molecules.

MHC Class II Molecules

MHC II molecules are both similar and different from MHC I molecules, as are their mechanisms of presentation. MHC II molecules are expressed on immune cells such as B cells, monocytes, macrophages, and dendritic cells and on epithelial cells following inflammatory signals, while MHC I molecules are expressed more ubiquitously. MHC II molecules on dendritic cells present antigen to naïve CD4⁺ T cells to activate them; later MHC II molecules participate in the interaction of B cells and macrophages with these specific CD4⁺ effector T cells [90]. This is a critical function, as exemplified by patients with deficits in MHC class II expression (bare lymphocyte syndrome), which results in extreme susceptibility to infections with a variety of microorganisms and death at a young age [91]. The structure of MHC class II resembles that of MHC class I and they are both polymorphic proteins (and thus transplantation antigens) [90,92]. Interestingly, *Gadus morhua* fish (Atlantic cod) lack MHC II but express an MHC I molecule containing endocytosis signals that effectively takes over MHC II function, illustrating the strong relationship and conserved function of these two MHC classes [93]. However, the natures of the



Trends in Immunology

Figure 6. A Simple Illustration of MHC II Antigen Presentation. MHC II proteins are made in the endoplasmic reticulum (ER) where they pair with a third chain, the invariant chain (Ii). Ii fills (through a peptide sequence called CLIP) the MHC II peptide-binding groove and allows efficient exit of MHC II from the ER. Ii also guides MHC II through the cell to a late endosomal compartment, MIIC. Ii is degraded by endosomal proteases, as are antigens taken up by endocytosis or phagocytosis. The CLIP segment is protected from destruction and exchanged for an antigenic peptide with the help of a dedicated chaperone called DM (HLA-DM in humans). Another chaperone expressed in a few immune cell types (immature B cells, some DC forms), called DO (HLA-DO), can compete for DM binding to MHC II and thereby affect the peptide repertoire on MHC II that is ultimately presented at the cell surface to CD4⁺ T cells.

presented peptides usually differ, as does the underlying biology of MHC class II antigen presentation.

How to Present the Outside World?

MHC class II presents peptide fragments that are generally larger than those presented by MHC class I, because the peptide-binding groove of MHC class II is open, allowing peptides to extend out of this site [94]. The MHC class II-associated peptides are derived from extracellular proteins and from self-proteins that are degraded in the endosomal pathway (Figure 6) [95]. MHC II molecules associate during their assembly in the ER with the invariant chain Ii, which acts as a pseudopeptide by filling the MHC class II peptide-binding groove and targets MHC II molecules into the endosomal pathway through its cytosolic dileucine motif [20,96]. In a compartment commonly named MIIC [97], MHC II molecules then meet antigenic fragments generated by resident proteases. For these peptides to bind MHC II molecules, the invariant chain has to be degraded by the same mix of proteases, especially cathepsin L and S [90]. This leaves an invariant chain fragment (called CLIP) inaccessible for proteases and remaining in the peptide-binding groove of MHC II molecules [98]. This CLIP fragment has to be exchanged for higher-affinity peptides with the help of a dedicated MHC class II-like chaperone called DM (in humans, HLA-DM) [99]. The structure of HLA-DM in association with MHC II (HLA-DR1) reveals that DM locally opens the groove to release low-affinity peptides such as CLIP. DM release from MHC

class II then locks the proper peptide fragments in the MHC class II peptide-binding groove [100]. After some period of residence in MIIC, MHC II molecules move to the plasma membrane either via vesicular transport or in the form of tubules [101–103]. Since the targeting information in the invariant chain has been removed after its degradation in MIIC, MHC II molecules can stably reside on the plasma membrane.

Variations in MHC Class II Presentation

As for MHC I molecules, there are various MHC II loci in most species (in humans there are three, named HLA-DR, HLA-DQ, and HLA-DP). Also, MHC II molecules are polymorphic (>3000 alleles known) [90] and their polymorphic amino acids similarly cluster in and around the peptide-binding groove, shaping the peptide-binding pockets. Consequently, different MHC II alleles bind different peptides by virtue of their different anchor residues [104]. This is likely to explain why certain MHC II alleles are strongly linked with defined autoimmune diseases [105] and immune responses against external environmental and food antigens [90]. This is well illustrated for the link between gluten sensitivity (celiac disease) and the MHC class II allele HLA-DQ2. HLA-DQ2 is able to present a peptide from gluten after deamination by tissue transglutaminase to activate CD4⁺ T cells and drives the disease [106]. The dominant antigens for autoimmune diseases are often unclear but there are some suggestions. For example, insulin has been suspected to mediate HLA-DR3/4-associated type 1 diabetes and myelin basic protein to participate in HLA-DR1 related multiple sclerosis [107,108]. One additional mechanism involves the presentation of an atypical conformation of a peptide [109] as the result of peptide loading of MHC II molecules in compartments that lack DM molecules, such as may occur in recycling endocytic compartments [110] or in the ER [111,112]. In the absence of DM function, the non-optimal peptide conformation bound to MHC II is not corrected. A different conformation of a self-peptide can be recognized as non-self by CD4⁺ T cells, which may drive the induction of autoantibodies [109]. However, the fact that many people with these MHC class II alleles never develop autoimmune diseases, and that for most of these conditions much less than half of identical twins are concordant for disease, indicates that epigenetic and other factors must also be involved and these are as yet unclear.

Besides genetic variation resulting in polymorphism of MHC II molecules, another variation in the life of MHC class II lies with the associated invariant chain, which is actually not so invariant. Indeed, there are multiple splice variants, including one (p44) that contains an additional protease inhibitor (cystatin) domain [90]. This variant can be expected to modify the protease activities involved in antigen preparation for MHC II molecules. The proteases (cathepsins in the endosomal pathway) also vary in the different MHC class II-expressing immune cells, as do natural inhibitors for these cathepsins (called cystatins) [90,96]. It is likely that antigens are degraded after denaturation (which involves reduction of their disulfide bonds by the enzyme GILT [113], and acidic pH [114]) by a swarm of proteases that will generate and destroy potential peptides for a defined MHC II allele.

Peptides are selected for presentation by MHC II with the help of the chaperone DM. DM can be further controlled by a dedicated co-chaperone called DO (HLA-DO in humans) that strongly pairs with DM [115]. DO associates with DM at the same interface as that occupied by MHC II molecules [100] and thus inhibits DM-assisted peptide loading of MHC II molecules unless the DM–DO complex enters very acidic endosomes along with MHC II molecules [116,117]. DO thus shapes the peptide repertoire by preventing peptide binding in earlier vesicular compartments. Deleting DO induces type 1 diabetes in mouse models and possibly other autoimmune diseases [118].

Other variations entail the transport and surface half-life of MHC II molecules. MHC II transport from MIIC to the plasma membrane is not constitutive but controlled in dendritic cells,

monocytes, and B cells. Activation of dendritic cells promotes MHC II transport to the cell surface and strongly enhances the half-life of these molecules on the plasma membrane [119,120]. As a result, activated dendritic cells have high numbers (around 2 million per cell) of MHC II molecules on the cell surface that continue to present antigens for long periods. However, there are other ways to control MHC II expression. For example, in human monocytes IL-10 increases the expression of a ubiquitin ligase membrane-associated RING-CH1 (MARCH-1) that ubiquitinates the tail of cell surface MHC II molecules initiating their rapid internalization and destruction [121]. Genome-wide analyses have identified many factors that control the complex process of control of MHC II expression and transport in dendritic cells [122]. Many pathogens, especially those residing in MHC II-containing phagosomes, also inhibit MHC II expression or peptide loading [123,124]. Such immune evasion by pathogens may be caused by manipulation of DM interactions with MHC II molecules, manipulation of pH levels, alteration of protein networks, or induction of MHC ubiquitination by MARCH homologs [125] or other processes required for optimal antigen presentation by MHC II molecules.

Concluding Remarks and Future Perspectives

The work of many laboratories over past decades has informed our understanding of the function of MHC class I and MHC class II molecules at the immunological, cell biological, genetic, and atomic level. Many of the major mechanisms of both the MHC I and the MHC II pathways are understood and we can look forward to an even more comprehensive understanding of antigen presentation mechanisms in the coming years. With such knowledge, we can look forward to better understanding how these processes help to maintain health and/or contribute to disease pathogenesis. We anticipate that there will be a strong potential to translate this knowledge into clinical medicine. One important area is in cancer immunotherapy. While the role of MHC class I and II antigen presentation in tumor immunology was initially restricted to curing cancer in mouse models, the recent development of checkpoint inhibitor antibodies has translated this into clinical responses in human melanoma, lung cancer, and other tumors [126]. Characterizing the defects in antigen presentation in such tumors might in theory help identify patients who can or cannot respond to such interventions. Moreover, if methods can be developed to reverse such defects, they might be able to improve the efficacy of these immunotherapies. Similarly, manipulating antigen presentation pathways might overcome the inhibition of antigen presentation induced by certain microbes and then help eliminate these chronic infections. The rules for antigen presentation by MHC I molecules can be used to predict neoepitopes detected by CD8⁺ T cells [127] and this information might be exploited to actively immunize cancer patients in conjunction with removal of checkpoint control and to monitor patient responses. Targeting antigens to dendritic cells and in particular into specific presentation pathways has the potential to generate more robust and effective kinds of responses to vaccines and immunotherapies. Antigen presentation might also be manipulated in the opposite way to dampen autoimmune diseases; for example, with tolerogenic peptides or tolerogenic antigen-presenting cells. Despite the remarkable advances in our knowledge about antigen presentation, the picture remains incomplete but should continue to improve. For example, forward genetics screens are uncovering unsuspected new components in the pathways [122,128]. Filling these gaps (see Outstanding Questions) should provide a higher-resolution understanding of the pathways and their contribution to disease pathogenesis as well as increase the opportunities to exploit these pathways to develop better immunotherapies to prevent and/or treat disease.

Acknowledgments

This work was supported by grants from NWO-TOP and an ERC Advanced Grant to J.N. and NIH grants RO1AI114495 and RO1AI110374 to K.L.R.

Supplemental Information

Supplemental information associated with this article can be found online at [doi:10.1016/j.it.2016.08.010](https://doi.org/10.1016/j.it.2016.08.010).

Outstanding Questions

While antigen presentation by MHC molecules has been intensively studied for 40 years, many aspects remain unclear. The following is only a partial list.

Why can peptides presented by MHC molecules not be more accurately predicted? What are we missing?

What are the many proteases involved in generating and destroying presented peptides and what are their relative contributions and specificities?

How do the antigen presentation pathways so successfully present a broad peptidome from very large numbers of different proteins of very different abundances and in the face of robust peptide destruction?

Are there mechanisms that protect peptides from destruction before they bind to MHC molecules?

The components of the MHC antigen presentation pathway have remarkable heterogeneity in expression and activity in different tissues. Does that lead to different presentation of self-antigens and contribute to autoimmune responses?

Why do we have three different proteasomes?

Why do we express only three MHC I and three MHC II locus products and not more to cover all possible peptides for presentation?

What is the major mechanism that generates DRIPS—inaccurate transcription, translation, folding, or assembly?

Is antigen cross-presentation the result of many different systems or are there dominant systems?

For cross-presentation, how are antigens transported from endosomes into the cytosol?

For cross-presentation, how are MHC I molecules kept stable during transport to and after arrival in endosomes?

Are there mechanisms that promote the loading of peptides in endosomes for cross-presentation?

References

- Flajnik, M.F. and Kasahara, M. (2010) Origin and evolution of the adaptive immune system: genetic events and selective pressures. *Nat. Rev. Genet.* 11, 47–59
- Brown, J.H. *et al.* (1993) Three-dimensional structure of the human class II histocompatibility antigen HLA-DR1. *Nature* 364, 33–39
- Bjorkman, P.J. *et al.* (1987) The foreign antigen binding site and T cell recognition regions of class I histocompatibility antigens. *Nature* 329, 512–518
- Zhang, P. *et al.* (2014) Crystal structure of the stress-inducible human heat shock protein 70 substrate-binding domain in complex with peptide substrate. *PLoS One* 9, e103518
- Dong, G. *et al.* (2009) Insights into MHC class I peptide loading from the structure of the tapasin-ERp57 thiol oxidoreductase heterodimer. *Immunity* 30, 21–32
- Teng, M.S. *et al.* (2002) A human TAPBP (TAPASIN)-related gene. *TAPBP-R. Eur. J. Immunol.* 32, 1059–1068
- Boyle, L.H. *et al.* (2013) Tapasin-related protein TAPBPR is an additional component of the MHC class I presentation pathway. *Proc. Natl. Acad. Sci. U.S.A.* 110, 3465–3470
- Mosyak, L. *et al.* (1998) The structure of HLA-DM, the peptide exchange catalyst that loads antigen onto class II MHC molecules during antigen presentation. *Immunity* 9, 377–383
- Fremont, D.H. *et al.* (1998) Crystal structure of mouse H2-M. *Immunity* 9, 385–393
- Guce, A.I. *et al.* (2013) HLA-DO acts as a substrate mimic to inhibit HLA-DM by a competitive mechanism. *Nat. Struct. Mol. Biol.* 20, 90–98
- Falk, K. *et al.* (1994) Pool sequencing of natural HLA-DR, DQ, and DP ligands reveals detailed peptide motifs, constraints of processing, and general rules. *Immunogenetics* 39, 230–242
- Falk, K. *et al.* (1991) Allele-specific motifs revealed by sequencing of self-peptides eluted from MHC molecules. *Nature* 351, 290–296
- Madden, D.R. *et al.* (1993) The antigenic identity of peptide–MHC complexes: a comparison of the conformations of five viral peptides presented by HLA-A2. *Cell* 75, 693–708
- Smith, K.J. *et al.* (1996) Bound water structure and polymorphic amino acids act together to allow the binding of different peptides to MHC class I HLA-B53. *Immunity* 4, 215–228
- Walz, S. *et al.* (2015) The antigenic landscape of multiple myeloma: mass spectrometry (re)defines targets for T-cell-based immunotherapy. *Blood* 126, 1203–1213
- Schmid, B.V. *et al.* (2010) Quantifying how MHC polymorphism prevents pathogens from adapting to the antigen presentation pathway. *Epidemics* 2, 99–108
- Chaix, R. *et al.* (2008) Is mate choice in humans MHC-dependent? *PLoS Genet.* 4, e1000184
- Siddle, H.V. *et al.* (2007) Characterization of major histocompatibility complex class I and class II genes from the Tasmanian devil (*Sarcophilus harrisii*). *Immunogenetics* 59, 753–760
- Siddle, H.V. *et al.* (2013) Reversible epigenetic down-regulation of MHC molecules by devil facial tumour disease illustrates immune escape by a contagious cancer. *Proc. Natl. Acad. Sci. U.S.A.* 110, 5103–5108
- Neefjes, J. *et al.* (2011) Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nat. Rev. Immunol.* 11, 823–836
- Michalek, M.T. *et al.* (1993) A role for the ubiquitin-dependent proteolytic pathway in MHC class I-restricted antigen presentation. *Nature* 363, 552–554
- Rock, K.L. *et al.* (1994) Inhibitors of the proteasome block the degradation of most cell proteins and the generation of peptides presented on MHC class I molecules. *Cell* 78, 761–771
- Reits, E. *et al.* (2003) Peptide diffusion, protection, and degradation in nuclear and cytoplasmic compartments before antigen presentation by MHC class I. *Immunity* 18, 97–108
- Cresswell, P. *et al.* (1999) The nature of the MHC class I peptide loading complex. *Immunol. Rev.* 172, 21–28
- Wearsch, P.A. and Cresswell, P. (2007) Selective loading of high-affinity peptides onto major histocompatibility complex class I molecules by the tapasin-ERp57 heterodimer. *Nat. Immunol.* 8, 873–881
- Garstka, M.A. *et al.* (2015) The first step of peptide selection in antigen presentation by MHC class I molecules. *Proc. Natl. Acad. Sci. U.S.A.* 112, 1505–1510
- Saveanu, L. *et al.* (2005) Concerted peptide trimming by human ERAP1 and ERAP2 aminopeptidase complexes in the endoplasmic reticulum. *Nat. Immunol.* 6, 689–697
- Saric, T. *et al.* (2002) An IFN- γ -induced aminopeptidase in the ER, ERAP1, trims precursors to MHC class I-presented peptides. *Nat. Immunol.* 3, 1169–1176
- Serwold, T. *et al.* (2002) ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* 419, 480–483
- Morozov, G.I. *et al.* (2016) Interaction of TAPBPR, a tapasin homolog, with MHC-I molecules promotes peptide editing. *Proc. Natl. Acad. Sci. U.S.A.* 113, E1006–E1015
- Hermann, C. *et al.* (2015) TAPBPR alters MHC class I peptide presentation by functioning as a peptide exchange catalyst. *Elife* 4, e09617
- Chang, S.C. *et al.* (2005) The ER aminopeptidase, ERAP1, trims precursors to lengths of MHC class I peptides by a “molecular ruler” mechanism. *Proc. Natl. Acad. Sci. U.S.A.* 102, 17107–17112
- York, I.A. *et al.* (2002) The ER aminopeptidase ERAP1 enhances or limits antigen presentation by trimming epitopes to 8–9 residues. *Nat. Immunol.* 3, 1177–1184
- Nguyen, T.T. *et al.* (2011) Structural basis for antigenic peptide precursor processing by the endoplasmic reticulum aminopeptidase ERAP1. *Nat. Struct. Mol. Biol.* 18, 604–613
- Roelse, J. *et al.* (1994) Trimming of TAP-translocated peptides in the endoplasmic reticulum and in the cytosol during recycling. *J. Exp. Med.* 180, 1591–1597
- Elliott, T. *et al.* (1991) Peptide-induced conformational change of the class I heavy chain. *Nature* 351, 402–406
- Schumacher, T.N. *et al.* (1990) Direct binding of peptide to empty MHC class I molecules on intact cells and *in vitro*. *Cell* 62, 563–567
- Kelly, A. *et al.* (1992) Assembly and function of the two ABC transporter proteins encoded in the human major histocompatibility complex. *Nature* 355, 641–644
- Neefjes, J.J. and Ploegh, H.L. (1988) Allele and locus-specific differences in cell surface expression and the association of HLA class I heavy chain with β 2-microglobulin: differential effects of inhibition of glycosylation on class I subunit association. *Eur. J. Immunol.* 18, 801–810
- Rammensee, H. *et al.* (1999) SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 50, 213–219
- Schubert, U. *et al.* (2000) Proteasome inhibition interferes with gag polyprotein processing, release, and maturation of HIV-1 and HIV-2. *Proc. Natl. Acad. Sci. U.S.A.* 97, 13057–13062
- Princiotta, M.F. *et al.* (2003) Quantitating protein synthesis, degradation, and endogenous antigen processing. *Immunity* 18, 343–354
- Reits, E.A. *et al.* (2000) The major substrates for TAP *in vivo* are derived from newly synthesized proteins. *Nature* 404, 774–778
- Yewdell, J.W. (2011) DRiPs solidify: progress in understanding endogenous MHC class I antigen processing. *Trends Immunol.* 32, 548–558
- Rock, K.L. *et al.* (2014) Re-examining class-I presentation and the DRiP hypothesis. *Trends Immunol.* 35, 144–152
- Monaco, J.J. and Nandi, D. (1995) The genetics of proteasomes and antigen processing. *Annu. Rev. Genet.* 29, 729–754
- Gaczynska, M. *et al.* (1993) γ -Interferon and expression of MHC genes regulate peptide hydrolysis by proteasomes. *Nature* 365, 264–267
- Groettrup, M. *et al.* (1996) A third interferon- γ -induced subunit exchange in the 20S proteasome. *Eur. J. Immunol.* 26, 863–869
- Toes, R.E. *et al.* (2001) Discrete cleavage motifs of constitutive and immunoproteasomes revealed by quantitative analysis of cleavage products. *J. Exp. Med.* 194, 1–12

Do dendritic cells or specific subsets of dendritic cells have unique components that promote cross-presentation?

How are hydrophobic peptides delivered in the peptide-binding groove of MHC I and MHC II molecules? Are there unique chaperones for such peptides?

Is there an MHC class II peptide-loading complex?

50. Kincaid, E.Z. *et al.* (2016) Specialized proteasome subunits have an essential role in the thymic selection of CD8⁺ T cells. *Nat. Immunol.* 17, 938–945
51. Ossendorp, F. *et al.* (1996) A single residue exchange within a viral CTL epitope alters proteasome-mediated degradation resulting in lack of antigen presentation. *Immunity* 5, 115–124
52. Tomaru, U. *et al.* (2009) Exclusive expression of proteasome subunit $\beta 5i$ in the human thymic cortex. *Blood* 113, 5186–5191
53. Yewdell, J.W. *et al.* (2003) Making sense of mass destruction: quantitating MHC class I antigen presentation. *Nat. Rev. Immunol.* 3, 952–961
54. Meyer, V.S. *et al.* (2009) Identification of natural MHC class II presented phosphopeptides and tumor-derived MHC class I phospholipids. *J. Proteome Res.* 8, 3666–3674
55. Haurum, J.S. *et al.* (1999) Presentation of cytosolic glycosylated peptides by human class I major histocompatibility complex molecules *in vivo*. *J. Exp. Med.* 190, 145–150
56. Petersen, J. *et al.* (2009) Post-translationally modified T cell epitopes: immune recognition and immunotherapy. *J. Mol. Med.* 87, 1045–1051
57. Gromme, M. *et al.* (1997) The rational design of TAP inhibitors using peptide substrate modifications and peptidomimetics. *Eur. J. Immunol.* 27, 898–904
58. Andersen, M.H. *et al.* (1999) Phosphorylated peptides can be transported by TAP molecules, presented by class I MHC molecules, and recognized by phosphopeptide-specific CTL. *J. Immunol.* 163, 3812–3818
59. Mohammed, F. *et al.* (2008) Phosphorylation-dependent interaction between antigenic peptides and MHC class I: a molecular basis for the presentation of transformed self. *Nat. Immunol.* 9, 1236–1243
60. Zarling, A.L. *et al.* (2000) Phosphorylated peptides are naturally processed and presented by major histocompatibility complex class I molecules *in vivo*. *J. Exp. Med.* 192, 1755–1762
61. Berkens, C.R. *et al.* (2015) Definition of proteasomal peptide splicing rules for high-efficiency spliced peptide presentation by MHC class I molecules. *J. Immunol.* 195, 4085–4095
62. Ploegh, H.L. (1998) Viral strategies of immune evasion. *Science* 280, 248–253
63. Ploegh, H.L. (1995) Trafficking and assembly of MHC molecules: how viruses elude the immune system. *Cold Spring Harb. Symp. Quant. Biol.* 60, 263–266
64. van Hall, T. *et al.* (2007) The Varicellovirus-encoded TAP inhibitor UL49.5 regulates the presentation of CTL epitopes by Qa-1b1. *J. Immunol.* 178, 657–662
65. Lichtenstein, D.L. and Wold, W.S. (2004) Experimental infections of humans with wild-type adenoviruses and with replication-competent adenovirus vectors: replication, safety, and transmission. *Cancer Gene Ther.* 11, 819–829
66. Ziegler, H. *et al.* (2000) The luminal part of the murine cytomegalovirus glycoprotein gp40 catalyzes the retention of MHC class I molecules. *EMBO J.* 19, 870–881
67. Mittal, D. *et al.* (2014) New insights into cancer immunoediting and its three component phases – elimination, equilibrium and escape. *Curr. Opin. Immunol.* 27, 16–25
68. Garcia-Lora, A. *et al.* (2003) MHC class I antigens, immune surveillance, and tumor immune escape. *J. Cell. Physiol.* 195, 346–355
69. Snyder, A. *et al.* (2014) Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* 371, 2189–2199
70. Smit, E.F. and Baas, P. (2016) Lung cancer in 2015: bypassing checkpoints, overcoming resistance, and honing in on new targets. *Nat. Rev. Clin. Oncol.* 13, 75–76
71. Saccheri, F. *et al.* (2010) Bacteria-induced gap junctions in tumors favor antigen cross-presentation and antitumor immunity. *Sci. Transl. Med.* 2, 44ra57
72. Nijssen, J. *et al.* (2005) Cross-presentation by intercellular peptide transfer through gap junctions. *Nature* 434, 83–88
73. Bevan, M.J. (1976) Cross-priming for a secondary cytotoxic response to minor H antigens with H-2 congenic cells which do not cross-react in the cytotoxic assay. *J. Exp. Med.* 143, 1283–1288
74. Rock, K.L. *et al.* (1990) Presentation of exogenous antigen with class I major histocompatibility complex molecules. *Science* 249, 918–921
75. Kurts, C. *et al.* (2010) Cross-priming in health and disease. *Nat. Rev. Immunol.* 10, 403–414
76. Amigorena, S. (2003) Y in X priming. *Nat. Immunol.* 4, 1047–1048
77. Segura, E. and Amigorena, S. (2015) Cross-presentation in mouse and human dendritic cells. *Adv. Immunol.* 127, 1–31
78. Guermonprez, P. *et al.* (2003) ER-phagosome fusion defines an MHC class I cross-presentation compartment in dendritic cells. *Nature* 425, 397–402
79. Zitvogel, L. *et al.* (2008) Immunological aspects of anticancer chemotherapy. *Bull. Acad. Natl. Med.* 192, 1469–1487
80. Kovacsics-Bankowski, M. *et al.* (1993) Efficient major histocompatibility complex class I presentation of exogenous antigen upon phagocytosis by macrophages. *Proc. Natl. Acad. Sci. U.S.A.* 90, 4942–4946
81. Matheoud, D. *et al.* (2010) Cross-presentation by dendritic cells from live cells induces protective immune responses *in vivo*. *Blood* 115, 4412–4420
82. Amigorena, S. (2002) Fc gamma receptors and cross-presentation in dendritic cells. *J. Exp. Med.* 195, F1–F3
83. Schuette, V. and Burgdorf, S. (2014) The ins-and-outs of endosomal antigens for cross-presentation. *Curr. Opin. Immunol.* 26, 63–68
84. Joffre, O.P. *et al.* (2012) Cross-presentation by dendritic cells. *Nat. Rev. Immunol.* 12, 557–569
85. Kovacsics-Bankowski, M. and Rock, K.L. (1995) A phagosome-to-cytosol pathway for exogenous antigens presented on MHC class I molecules. *Science* 267, 243–246
86. Gromme, M. *et al.* (1999) Recycling MHC class I molecules and endosomal peptide loading. *Proc. Natl. Acad. Sci. U.S.A.* 96, 10326–10331
87. Dorsey, B.D. *et al.* (2008) Discovery of a potent, selective, and orally active proteasome inhibitor for the treatment of cancer. *J. Med. Chem.* 51, 1068–1072
88. Shen, L. *et al.* (2004) Important role of cathepsin S in generating peptides for TAP-independent MHC class I crosspresentation *in vivo*. *Immunity* 21, 155–165
89. Nair-Gupta, P. *et al.* (2014) TLR signals induce phagosomal MHC-I delivery from the endosomal recycling compartment to allow cross-presentation. *Cell* 158, 506–521
90. Unanue, E.R. *et al.* (2016) Variations in MHC class II antigen processing and presentation in health and disease. *Annu. Rev. Immunol.* 34, 265–297
91. Waldburger, J.M. *et al.* (2000) Lessons from the bare lymphocyte syndrome: molecular mechanisms regulating MHC class II expression. *Immunol. Rev.* 178, 148–165
92. Jones, E.Y. (1997) MHC class I and class II structures. *Curr. Opin. Immunol.* 9, 75–79
93. Malmstrom, M. *et al.* (2013) Unraveling the evolution of the Atlantic cod's (*Gadus morhua* L.) alternative immune strategy. *PLoS One* 8, e74004
94. Stern, L.J. *et al.* (1994) Crystal structure of the human class II MHC protein HLA-DR1 complexed with an influenza virus peptide. *Nature* 368, 215–221
95. Suri, A. *et al.* (2006) The wide diversity and complexity of peptides bound to class II MHC molecules. *Curr. Opin. Immunol.* 18, 70–77
96. Cresswell, P. and Roche, P.A. (2014) Invariant chain-MHC class II complexes: always odd and never invariant. *Immunol. Cell Biol.* 92, 471–472
97. Neeffjes, J. (1999) CIIV, MHC and other compartments for MHC class II loading. *Eur. J. Immunol.* 29, 1421–1425
98. Ghosh, P. *et al.* (1995) The structure of an intermediate in class II MHC maturation: CLIP bound to HLA-DR3. *Nature* 378, 457–462

99. Denzin, L.K. and Cresswell, P. (1995) HLA-DM induces CLIP dissociation from MHC class II alpha beta dimers and facilitates peptide loading. *Cell* 82, 155–165
100. Pos, W. *et al.* (2012) Crystal structure of the HLA-DM–HLA-DR1 complex defines mechanisms for rapid peptide selection. *Cell* 151, 1557–1568
101. Wubbolts, R. *et al.* (1996) Direct vesicular transport of MHC class II molecules from lysosomal structures to the cell surface. *J. Cell Biol.* 135, 611–622
102. Boes, M. *et al.* (2002) T-cell engagement of dendritic cells rapidly rearranges MHC class II transport. *Nature* 418, 983–988
103. Kleijmeer, M. *et al.* (2001) Reorganization of multivesicular bodies regulates MHC class II antigen presentation by dendritic cells. *J. Cell Biol.* 155, 53–63
104. Nielsen, M. *et al.* (2010) NetMHCIIpan-2.0 – improved pan-specific HLA-DR predictions using a novel concurrent alignment and weight optimization training procedure. *Immunome Res.* 6, 9
105. Fernando, M.M. *et al.* (2008) Defining the role of the MHC in autoimmunity: a review and pooled analysis. *PLoS Genet.* 4, e1000024
106. Solid, L.M. and Jabri, B. (2011) Celiac disease and transglutaminase 2: a model for posttranslational modification of antigens and HLA association in the pathogenesis of autoimmune disorders. *Curr. Opin. Immunol.* 23, 732–738
107. Steck, A.K. and Rewers, M.J. (2011) Genetics of type 1 diabetes. *Clin. Chem.* 57, 176–185
108. Pette, M. *et al.* (1990) Myelin autoreactivity in multiple sclerosis: recognition of myelin basic protein in the context of HLA-DR2 products by T lymphocytes of multiple-sclerosis patients and healthy donors. *Proc. Natl. Acad. Sci. U.S.A.* 87, 7968–7972
109. Wan, X. *et al.* (2016) Class-switched anti-insulin antibodies originate from unconventional antigen presentation in multiple lymphoid sites. *J. Exp. Med.* 213, 967–978
110. Sinnathamby, G. and Eisenlohr, L.C. (2003) Presentation by recycling MHC class II molecules of an influenza hemagglutinin-derived epitope that is revealed in the early endosome by acidification. *J. Immunol.* 170, 3504–3513
111. Tewari, M.K. *et al.* (2005) A cytosolic pathway for MHC class II-restricted antigen processing that is proteasome and TAP dependent. *Nat. Immunol.* 6, 287–294
112. van Luijn, M.M. *et al.* (2010) Alternative li-independent antigen-processing pathway in leukemic blasts involves TAP-dependent peptide loading of HLA class II complexes. *Cancer Immunol. Immunother.* 59, 1825–1838
113. Maric, M. *et al.* (2001) Defective antigen processing in GILT-free mice. *Science* 294, 1361–1365
114. Ziegler, H.K. and Unanue, E.R. (1982) Decrease in macrophage antigen catabolism caused by ammonia and chloroquine is associated with inhibition of antigen presentation to T cells. *Proc. Natl. Acad. Sci. U.S.A.* 79, 175–178
115. Liljedahl, M. *et al.* (1996) HLA-DO is a lysosomal resident which requires association with HLA-DM for efficient intracellular transport. *EMBO J.* 15, 4817–4824
116. Denzin, L.K. *et al.* (1997) Negative regulation by HLA-DO of MHC class II-restricted antigen processing. *Science* 278, 106–109
117. van Ham, S.M. *et al.* (1997) HLA-DO is a negative modulator of HLA-DM-mediated MHC class II peptide loading. *Curr. Biol.* 7, 950–957
118. Yi, W. *et al.* (2010) Targeted regulation of self-peptide presentation prevents type I diabetes in mice without disrupting general immunocompetence. *J. Clin. Invest.* 120, 1324–1336
119. Cella, M. *et al.* (1997) Inflammatory stimuli induce accumulation of MHC class II complexes on dendritic cells. *Nature* 388, 782–787
120. Pierre, P. *et al.* (1997) Developmental regulation of MHC class II transport in mouse dendritic cells. *Nature* 388, 787–792
121. Thibodeau, J. *et al.* (2008) Interleukin-10-induced MARCH1 mediates intracellular sequestration of MHC class II in monocytes. *Eur. J. Immunol.* 38, 1225–1230
122. Paul, P. *et al.* (2011) A genome-wide multidimensional RNAi screen reveals pathways controlling MHC class II antigen presentation. *Cell* 145, 268–283
123. Mitchell, E.K. *et al.* (2004) Inhibition of cell surface MHC class II expression by *Salmonella*. *Eur. J. Immunol.* 34, 2559–2567
124. Zwart, W. *et al.* (2005) Spatial separation of HLA-DM/HLA-DR interactions within MHC and phagosome-induced immune escape. *Immunity* 22, 221–233
125. van Niel, G. *et al.* (2006) Dendritic cells regulate exposure of MHC class II at their plasma membrane by oligoubiquitination. *Immunity* 25, 885–894
126. Pennock, G.K. and Chow, L.Q. (2015) The evolving role of immune checkpoint inhibitors in cancer treatment. *Oncologist* 20, 812–822
127. Schumacher, T.N. and Schreiber, R.D. (2015) Neoantigens in cancer immunotherapy. *Science* 348, 69–74
128. Jongsma, M.L. *et al.* (2016) An ER-associated pathway defines endosomal architecture for controlled cargo transport. *Cell* 166, 152–166
129. Saveanu, L. *et al.* (2009) IRAP identifies an endosomal compartment required for MHC class I cross-presentation. *Science* 325, 213–217